

RESCUE[®]

CONCENTRATE

ONE STEP DISINFECTANT CLEANER & DEODORIZER

Reference Sheet

For Veterinary Use

Manufactured by:
Virox Technologies Inc.
2770 Coventry Road
Oakville, ON Canada L6H 6R1
viroxanimalhealth.com



When used as directed, this product is highly effective against a wide variety of pathogenic microorganisms including bacteria, antibiotic-resistant bacteria, *viruses, fungi, mold and mildew. At a 1:16 dilution (8 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load and 5 minute contact time, unless otherwise noted, this product kills the following on hard, non-porous inanimate surfaces:

BACTERICIDAL: 5 MINUTES @ 1:16

Acinetobacter baumannii
Escherichia coli O157:H7
Klebsiella pneumoniae
Listeria monocytogenes
Pseudomonas aeruginosa
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Shigella dysenteriae
Staphylococcus aureus
Streptococcus pyogenes

ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY:

Enterococcus faecium (Vancomycin Resistant (VRE))
Escherichia coli (Extended Spectrum Beta-lactamase resistance (ESBL))
Klebsiella pneumoniae (Carbapenem-resistant (KPC))
Staphylococcus aureus (CA-MRSA) (Genotype USA300)
Staphylococcus aureus (CA-MRSA) (Genotype USA400)
Staphylococcus aureus (MRSA)
Staphylococcus epidermidis (MRSE)
Streptococcus pneumoniae (Penicillin Resistant (PRSP))

*VIRUCIDAL: 5 MINUTES @ 1:64

At 1:16-1:64 dilution (2-8 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load and 5 minute contact time, unless otherwise noted, this product kills the following on hard, non-porous inanimate surfaces:

Adenovirus type 8
Herpes simplex Type 2
Human Coronavirus
Influenza Type A₂
Norovirus (Feline Calicivirus as the surrogate)
Parainfluenza Type 3
Poliovirus Type 1
Respiratory Syncytial Virus
Rhinovirus Type 37, Strain 151-1
Rotavirus
Vaccinia Virus (smallpox vaccine virus)
Kills *Hepatitis B Virus (HBV) & *Hepatitis C Virus (HCV) when used as directed on hard, non-porous inanimate surfaces with a 5 minute contact time.
Kills *HIV-1 (AIDS virus) (HTLV-III_B) when used as directed on hard, non-porous inanimate surfaces with a 1 minute contact time.

*This product has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.

VETERINARY *VIRUSES

Avian Influenza A (H3N2)
Canine Distemper virus
Canine Adenovirus
Canine Coronavirus
Canine Parainfluenza
Feline Infectious Peritonitis
Feline Picornavirus
Feline Rhinotracheitis virus
Pseudorabies virus
Equine Herpes virus type 1
Equine Influenza A (H3N8) virus
Porcine Respiratory & Reproductive Syndrome (PRRS) virus
Porcine Epidemic Diarrhea virus

*Canine Parvovirus: Kills *Canine Parvovirus when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 minute contact time.

*Canine Parvovirus: Kills *Canine Parvovirus when used as directed on hard, non-porous inanimate surfaces at a 1:32 dilution (4 oz. per gallon of water), and in the presence of a 5% serum load and 10 minute contact time.

*Minute Virus of Mice: Kills *Minute Virus of Mice when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 minute contact time.

FUNGICIDAL: 5 MINUTES @ 1:16

This product is fungicidal when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 minute contact time at 20°C (68°F).

Trichophyton mentagrophytes
Microsporum canis

MALODORS

Malodors – reduces odors and odor-causing bacteria on hard non-porous surfaces in restroom areas, behind and under sinks and counters, and storage areas and other places where bacterial growth can cause malodors, when used as directed at a 1:16 dilution (8 oz. of product per gallon of water), in the presence of 200 ppm hard water and a 30 seconds contact time.

MOLD/MILDEW

Mildewcidal Activity: Kills the growth of mold and mildew when diluted at 1:16 in 200 ppm hard water at a 10 minute contact time: *Aspergillus niger* and the odors caused by them when applied to hard, non-porous environmental surfaces.

Mildewstatic Activity: Controls and prevents the growth of mold and mildew when diluted at 1:16 in 200 ppm hard water at a 10 minute contact time: *Aspergillus niger* and the odors caused by them when applied to hard, non-porous environmental surfaces.

Si no puede leer en inglés, pregunte a su supervisor sobre las instrucciones de uso apropiadas antes de trabajar con este producto.

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER: 30 SECONDS @ 1:16

Sanitizing: When used as a non-food contact sanitizer at a 1:16 dilution (8 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load, this product reduces the following bacteria by 99.9% with a 30 second contact time on hard non-porous inanimate surfaces:

Enterobacter aerogenes

Enterococcus faecalis (Vancomycin Resistant (VRE))

Escherichia coli (Extended Spectrum Beta-lactamase resistance (ESBL))

Listeria monocytogenes

Pseudomonas aeruginosa

Salmonella enterica (formerly known as *Salmonella choleraesuis*)

Staphylococcus aureus

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER: 3 MINUTES @ 1:128

Sanitizing: When used as a non-food contact sanitizer at a 1:128 dilution (1 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load, this product reduces the following bacteria by 99.9% with a 3 minute contact time on hard non-porous inanimate surfaces:

Escherichia coli

Enterococcus faecalis

Klebsiella pneumoniae

Listeria monocytogenes

Pseudomonas aeruginosa

Salmonella enterica (formerly known as *Salmonella choleraesuis*)

Staphylococcus aureus

BACTERICIDAL STABILITY OF USE-SOLUTION

Tests show this product as a disinfectant, when diluted at 1:16 dilution (8 oz. of product per gallon of water) in 200 ppm hard water, in the presence of 5% serum, remains effective against *Staphylococcus aureus*, *Salmonella enterica* (formerly known as *Salmonella choleraesuis*) and *Pseudomonas aeruginosa* for up to 90 days in storage as long as the bottle remains sealed.

If the use-solution becomes visibly dirty or contaminated, it must be discarded and a fresh product prepared. Always use clean, dry containers when diluting this product. Treated surfaces must remain wet for 5 minutes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

*This product kills HBV, HCV and HIV-1 on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in health care settings and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Hepatitis B Virus (HBV), Hepatitis C Virus (HCV) and Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HBV, HCV and HIV-1 OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood and body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1, and for 5 minutes to kill HBV and HCV.

Disposal of Infectious Material: Cleaning materials used that may contain blood and other body fluids must be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

For use as a *Virucide: Dilute at 2 – 8 oz. of product per gallon of water (1:16 – 1:64). **1.** Pre-clean heavily soiled areas. **2.** Apply Use Solution until thoroughly wet. **3.** Let stand for 5 minutes. **4.** Wipe surfaces and let air dry.

To Kill *Minute Virus of Mice: Dilute at 8.0 oz. of product per gallon of water (Dilute 1:16). **1.** Pre-clean heavily soiled areas. **2.** Apply Use Solution until thoroughly wet. **3.** Let stand for 5 minutes. **4.** Wipe surfaces and let air dry.

To Kill *Canine Parvovirus: Dilute at 4.0 oz. of product per gallon of water (Dilute 1:16). **1.** Pre-clean heavily soiled areas. **2.** Apply Use Solution until thoroughly wet. **3.** Let stand for 10 minutes. **4.** Wipe surfaces and let air dry.

To Kill Mold and Mildew: Dilute at 8.0 oz. of product per gallon of water (1:16). Pre-clean surfaces. Apply Use Solution to hard, non-porous environmental surfaces. Allow surfaces to remain wet for 10 minutes. Wipe surfaces and let air dry.

To Control Mold and Mildew: Dilute at 8.0 oz. of product per gallon of water (1:16). Apply Use Solution to pre-cleaned hard, non-porous environmental surfaces. Wipe surfaces and let air dry. Repeat application weekly or when growth reappears.

To Kill Fungi: Dilute at 8.0 oz. of product per gallon of water (1:16). Pre-clean heavily soiled areas. Apply Use Solution to hard, non-porous environmental surfaces. Allow surface to remain wet for 5 minutes. Wipe surfaces and let air dry.

To Sanitize Non-Food Contact Surfaces: Dilute at 8.0 oz. of product per gallon of water (Dilute 1:16). **1.** Pre-clean heavily soiled hard non-porous surfaces. **2.** Apply Use Solution until thoroughly wet. **3.** Let stand for 30 seconds. **4.** Wipe surfaces and let air dry. **5.** Not for use on food contact surfaces or on food preparation areas.

To Sanitize Non-Food Contact Surfaces: Dilute at 1.0 oz. of product per gallon of water (Dilute 1:128). **1.** Pre-clean heavily soiled hard non-porous surfaces. **2.** Apply Use Solution until thoroughly wet. **3.** Let stand for 3 minutes. **4.** Wipe surfaces and let air dry. **5.** Not for use on food contact surfaces or on food preparation areas.

For Treatment of Animal Housing Facilities: Dilute at 8.0 oz. of product per gallon of water (1:16).

1. Remove all animals and feeds from areas being treated.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities occupied or traversed by animals.
3. Empty or cover all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
5. Apply fresh Use Solution to floors, walls, cages and other washable hard, non-porous environmental surfaces. For smaller surfaces, use a trigger spray bottle to spray all surfaces with solution until wet. To disinfect, all surfaces must remain wet for 5 minutes.
6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings and other closed spaces. Do not house animals or re-employ equipment until product has dried.
8. For disinfection of feed racks, troughs, automatic feeders, fountains and watering appliances scrub with use-solution, let stand 5 minutes. Then thoroughly scrub all treated surfaces with soap or detergent and rinse with potable water before reuse.